

# Drug Information Sheet("Kusuri-no-Shiori")

External

Revised: 05/2025

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:**DORMOLOL COMBINATION OPHTHALMIC SOLUTION  
"WAKAMOTO"

**Active ingredient:**Dorzolamide hydrochloride  
Timolol maleate

**Dosage form:**limpid and colorless, slightly viscous, ophthalmic solution, 5mL, (cap)  
orange

**Imprint or print on wrapping:**



## Effects of this medicine

This medicine lowers pressure in the eye by reducing production of aqueous humor by inhibiting carbonic anhydrase and blocking beta-receptor.

It is usually used to treat glaucoma or ocular hypertension.

## The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.  
If you have: bronchial asthma/history of bronchial asthma, bronchospasm, chronic obstructive pulmonary disease, heart failure, sinus bradycardia, atrioventricular block, cardiogenic shock, renal disorder, right-sided heart failure secondary to pulmonary hypertension, congestive heart failure, diabetic ketoacidosis, metabolic acidosis, diabetes, liver dysfunction, or history of intraocular surgery.
- If you wear contact lenses.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, apply 1 drop at a time, twice a day. Strictly follow the instructions.
- Make sure not to let the tip of the container touch any part of your eye.
- After applying, keep your eyes closed without blinking for a while (1 to 5 minutes).
- If this medicine gets on your eyelid, wipe it off immediately.
- If you need to use other eye drops at the same time, wait more than 5 minutes.
- If you miss a dose, apply the missed dose as soon as possible. However, if it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

## Precautions while taking this medicine

## Possible adverse reactions to this medicine

The most commonly reported adverse reactions include stinging eyes (smarting/burning sensation/foreign body sensation/lacrimation/pain/itchy eyes). If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

- foreign body sensation in eyes, eye pain, bloodshot eyes [ocular pemphigoid]
- shortness of breath, breathing difficulty, wheezing (whistling sound) [bronchospasm, respiratory distress, respiratory failure]
- respiratory distress, loss of consciousness, dizziness on standing up [heart block, congestive heart failure, cerebral ischemia, cardiac arrest, cerebrovascular disorder]
- fever, facial erythema, general malaise [systemic lupus erythematosus]
- fever, redness/erosion of mucosa, erythema [Stevens-Jonson syndrome, toxic epidermal necrolysis]

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

## Storage conditions and other information

- Keep out of the reach of children. Store away from light, heat and moisture. Store at room temperature (1 to 30 degrees Celsius) in the accompanying bag.

•Discard the remainder that has passed 1 month after opening without storing it. If you do not know how to dispose of it, please consult the pharmacy or medical institution where you received it. Do not give it to others.

**For healthcare professional use only**      /      /

For further information, talk to your doctor or pharmacist.